

2-year-old male • Left leg • 6% TBSA • PermeaDerm® Sheet

Day 0



Wound assessment

Day 7



Secondary adherence completed

Day 14



100% Re-epithelialization

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.





Day 0



Wound assessment

Day 7



Secondary adherence completed

Day 14



100% Re-epithelialization

Day 28



Continuous improvement in appearance

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.



Case Report: Deep Partial-Thickness Burn





Wound assessment



Secondary adherence completed




80% Re-epithelialization

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.



Case Report: Deep Partial-Thickness Burn





Secondary adherence completed

100% Re-epithelialization

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.







Wound assessment



Secondary adherence completed



100% Re-epithelialization

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.





1-year-old female • Left Arm • 1% TBSA • PermeaDerm® Sheet

Day 0

Wound assessment



Secondary adherence completed



100% Re-epithelialization

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.



10-year-old female • Right Forearm • 2% TBSA • PermeaDerm® Sheet

Day 0



Wound assessment

Day 7



Secondary adherence completed

Day 14



100% Re-epithelialization

6 Months



Optimal outcome with no scar

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.





23-year-old female • Right Foot • 1% TBSA • PermeaDerm® Sheet

Day 0



Wound assessment

Day 9



Secondary adherence completed

Day 16



100% Re-epithelialization

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.





Day 0



Wound assessment

Day 6



Secondary adherence completed

Day 14



100% Re-epithelialization

Day 21



Continuous improvement in appearance

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.





Day 0



Wound assessment

Day 6



Secondary adherence completed

Day 14



100% Re-epithelialization

Day 21



Optimal outcome with no scar

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.





15-year-old female • Right Breast • 1% TBSA • PermeaDerm® Sheet

Day 0



Wound assessment

Day 7




Secondary adherence completed

Day 14



90% Re-epithelialization

Day 21



100% Re-epithelialization

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.





59-year-old male • Left Hand • 2% TBSA • PermeaDerm® Sheet

Day 0



Wound assessment

Day 7



Secondary adherence completed

Day 14



98% Re-epithelialization

Day 21



Continuous improvement in appearance

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.



Case Report: Deep Partial-Thickness Burn




12-year-old female • Right Leg • 6% TBSA • PermeaDerm® Sheet

Day 0



Wound assessment

Day 7



Secondary adherence completed

Day 14



100% Re-epithelialization

6 Months



Optimal outcome with no scar

Consult the Instructions for Use at avitamedical.com for complete Safety Information prior to using PermeaDerm.

